



**University of California**  
Agriculture and Natural Resources




*Making a Difference  
for California*





HEALTH

**University of California**  
Agriculture and Natural Resources



*Making a Difference  
for California*



# **Large scale, low cost restoration of *Stipa pulchra* grasslands using herbicides.**

**Carl Bell, Marti Witter, and John Eckhoff**

**University of California  
Cooperative Extension, University  
of California Riverside, US  
National Park Service, CA  
Department of Fish and Game**



**University of California**  
Agriculture and Natural Resources

**Invasive Plants**



## Why Herbicides?

- ❖ Low cost
- ❖ Effective
- ❖ Can be used early in the rainy season
  - ❖ Kill the weeds
  - ❖ Let the natives have the water
- ❖ Environmental pros and cons equivalent to other methods such as burning or grazing

**University of California**  
Agriculture and Natural Resources



San Diego  
Sites –  
CA DFG  
Rancho  
Jamul  
Ecological  
Reserve

Los Angeles Co. sites  
-US NPS Santa  
Monica Mountains  
National Recreation  
Area, Cheeseboro  
Canyon

- ❖ In an area with an existing, but sparse native stand of purple needlegrass;
  - ❖ Can we find post-emergence herbicides that are safe to apply **broadcast** over the native grass?
  - ❖ Will these herbicide treatments increase the cover and vigor of the needlegrass?

© 2011 Google  
© 2011 Europa Technologies  
© 2011 INEGI

©2010 Google™


Imagery Date: 8/23/2010 1994

32°39'42.36" N 116°51'39.32" W elev 266 m

Eye alt 371 m

HEALTH

University of California  
Agriculture and Natural Resources





## Herbicide Selectivity can be based upon:

- ❖ Genetically different susceptibility
  - ❖ At the Class to Species level
  - ❖ Life history (annual vs perennial)
  - ❖ Or unique physiology
- ❖ Rate = amount of herbicide applied
- ❖ Plant phenology
- ❖ Season

**University of California**  
Agriculture and Natural Resources



- Herbicides applied in spring to established purple needlegrass, annually for 3 years.
- Herbicide treatments include:
  - Fluazifop-butyl (Fusilade)
  - Clethodim (Envoy)
  - Glyphosate (Roundup)
  - Pelargonic acid (Scythe)
  - Imazapic (Plateau)
  - Trifluralin (Preen granules)
  - Aminopyralid (Milestone)
  - Triclopyr (Garlon)



# SMMNRA Research

- Herbicide applied Jan. 2007 and Feb. 2009
- Plots sampled about 4 MAT for biomass, basal diameter, inflorescences per plant, and visual injury
- No significant difference ( $p=0.19$ ) between treatments and the UTC for biomass
- No apparent injury related to treatment



# RJER Results

Treatment	Cover		Grams/plant		Basal diameter (cms)	
	site 1	site 2	site 1	site 2	Site 1	Site 2
Fusilade + Garlon LR	23.3	10.5	48.0	19.8	89	94
Fusilade + Garlon HR	30.0	18.8	56.0	21.8	72	106
Glyphosate LR	6.3	15	12.5	17.5	37	76
Glyphosate HR	0.3	9.5	0.8	12.5	7	80
UTC	0.5	6.3	9.0	8.5	23	67


**University of California**  
Agriculture and Natural Resources







**University of California**  
Agriculture and Natural Resources



*Making a Difference  
for California*



- ❖ There are no silver bullets
  - ❖ Fusilade alone was not sufficient
    - ❖ *Vulpia myuros* is resistant to Fusilade and all other grass herbicides
  - ❖ Glyphosate was safe and effective in some cases but not all
  - ❖ Fusilade plus Garlon or Milestone worked well, but;
    - ❖ Milestone and Garlon damage geophytes
- ❖ Regardless, some treatments increased needlegrass cover and vigor significantly
  - ❖ Eliminating weeds early in spring benefits natives
  - ❖ Treatments are low cost and effective
  - ❖ Integrated weed management systems can be developed

